

THE ℓ -ADIC K -THEORY OF A p -LOCAL FIELD

GRACE K. LYO (MASSACHUSETTS INSTITUTE OF TECHNOLOGY)

ABSTRACT. We verify a special case of a conjecture of G. Carlsson that describes the ℓ -adic K -theory of a field F of characteristic prime to ℓ in terms of the representation theory of the absolute Galois group G_F . This conjecture is known to hold in two cases; in this article we examine the second case, in which the field in question is the field of Laurent series over a finite field $\mathbb{F}_p((x))$.

1. INTRODUCTION

We verify a special case of a conjecture of G. Carlsson that describes the ℓ -adic K -theory of a field F of characteristic prime to ℓ in terms of the representation theory of the absolute Galois group G_F . This conjecture employs a completion construction of Carlsson's called the *derived completion* [Car07a], which generalizes Bousfield and Kan's ℓ -adic completion [BK72] to the categories of ring and module spectra. Throughout this article, we will work in the category of $\mathbb{S}$ -algebras (which we will refer to as simply “ring spectra”) from [EKMM97]. If A is a commutative ring spectrum, M is an A -module, and $\gamma: A \rightarrow B$ is a map of commutative ring spectra, then we may take the derived completion of M with respect to γ ; we will write $M_\gamma^\wedge$ or $M_B^\wedge$ to denote this completion.

We will be taking derived completions of K -theory spectra of categories of representations. For any field E equipped with the action of a group G , denote by $\text{Rep}_E[G]$ the category of finite dimensional continuous E -linear representations of G and by $\text{Rep}_E\langle G \rangle$ the analogous category of E -semilinear representations. We define an E -semilinear representation of G to be an E -vector space V with a G -action satisfying the relation

$$g(a \cdot v) = {}^g a \cdot g(v),$$

where $g \in G$, $a \in E$, $v \in V$, ${}^g a$ is the image of a under the action by g on E , and $g(v)$ is the image of v under the action by g on V . It was shown in [EM06] that the spectra $K\text{Rep}_E\langle G \rangle$ are commutative $\mathbb{S}$ -algebras.

Denote by $\mathfrak{F}_\ell$ the symmetric monoidal category with ℓ objects, only identity morphisms whose monoid structure is determined by the additive structure of $\mathbb{F}_\ell$. All derived completions will be taken with respect to a map

$$(1) \quad \alpha_\ell: K\text{Rep}_E\langle G \rangle \longrightarrow K\mathbb{F}_\ell \cong H\mathfrak{F}_\ell,$$

for a prime ℓ different from the characteristic of E . This map is induced by the augmentation functor from $\text{Rep}_E\langle G \rangle$ to $\mathfrak{F}_\ell$, which sends a vector space to its E -dimension modulo ℓ .

Let F be a field and k an algebraically closed subfield of $\overline{F}$. Since k is algebraically closed, it is preserved by the absolute Galois group G_F and hence there is a category $\text{Rep}_k\langle G_F \rangle$. Define $\mathcal{E}_k^{\overline{F}}: \text{Rep}_k\langle G_F \rangle \longrightarrow \text{Rep}_{\overline{F}}\langle G_F \rangle$ to be the extension of scalars functor, where the action by G_F on k and on $\overline{F}$ is the Galois action

and where the action by G_F on $\mathcal{E}_k^{\overline{F}}(V)$ is the obvious one. Denote by $KF_\ell^\wedge$ the Bousfield–Kan ℓ -adic completion.

Conjecture (Carlsson). *For all primes ℓ different from the characteristic of F , the induced map on completions (which we also denote $\mathcal{E}_k^{\overline{F}}$)*

$$\mathcal{E}_k^{\overline{F}} : (K\mathrm{Rep}_k\langle G_F \rangle)_{\alpha_\ell}^\wedge \longrightarrow (K\mathrm{Rep}_{\overline{F}}\langle G_F \rangle)_{\alpha_\ell}^\wedge \simeq (KF)_\ell^\wedge$$

is a weak equivalence.

Note that the weak equivalence on the right is a result of two facts. First, the category $\mathrm{Rep}_E\langle G \rangle$ is equivalent to the category $\mathrm{Vect}(E^G)$ of finite dimensional E^G -vector spaces that arises whenever G acts faithfully on E . Second, the derived completion $(KF)_{\alpha_\ell}^\wedge$ becomes the Bousfield–Kan ℓ -adic completion when ℓ is prime to the characteristic of F [Car07b].

In [Car07a], Carlsson proved that this conjecture holds when F is a characteristic zero field containing an algebraically closed subfield k , provided G_F is abelian.

Fix primes $\ell \neq p$, and for the remainder of this article, let $k := \overline{\mathbb{F}}_p$ and define the field $\mathcal{F}$ to be the compositum of all p and ℓ -prime extensions of the field of Laurent series $\mathbb{F}_p((x))$. Denote by $\mathcal{F}^t$ the maximal tame extension of $\mathcal{F}$ and $G_{\mathcal{F}}^t$ the tame Galois group. Our main goal is to prove the following theorem.

Theorem 5.4. *The map*

$$(2) \quad \mathcal{E}_k^{\mathcal{F}^t} : (K\mathrm{Rep}_k\langle G_{\mathcal{F}}^t \rangle)_{\alpha_\ell}^\wedge \xrightarrow{\simeq} (K\mathrm{Rep}_{\mathcal{F}^t}\langle G_{\mathcal{F}}^t \rangle)_{\alpha_\ell}^\wedge \simeq (K\mathcal{F})_\ell^\wedge,$$

where $\mathcal{F}^t$ is the maximal tame extension of $\mathcal{F}$, is a weak equivalence.

For simplicity, we will write $\mathcal{E} = \mathcal{E}_k^{\mathcal{F}^t}$.

Note that in this theorem we use the tame Galois group $G_{\mathcal{F}}^t$, which we denote by $\mathcal{G}$ throughout this article, and the maximal tame extension $\mathcal{F}^t$, whereas the conjecture regards the absolute Galois group and the separable closure.

This substitution is justified by the fact that if P is a normal p -power subgroup of a profinite group G , and G acts on a characteristic p field E , then there is a weak equivalence

$$K\mathrm{Rep}_E\langle G/P \rangle \simeq K\mathrm{Rep}_E\langle G \rangle.$$

Combining this with (2) gives the conjectured equivalence $\mathcal{E}_k^{\overline{F}}$.

The first step in our strategy towards proving this theorem will be to compute the K -groups $K_*\mathrm{Rep}_k\langle \mathcal{G} \rangle$, (Section 4). We plug this data in to the derived completion’s algebraic to geometric spectral sequence, which converges to $\pi_*((K\mathrm{Rep}_k\langle \mathcal{G} \rangle)_{\alpha_\ell}^\wedge)$. In order to do this, we use results from “Semisimple skew group rings and their modules” [LR07], which is joint work with K. Ribet. We summarize the relevant portions of this article in Section 3. Applying these results yields the following propositions about the homotopy groups of $K\mathrm{Rep}_k\langle \mathcal{G} \rangle$.

Proposition 4.4. *The endomorphism rings of the irreducible representations in $\mathrm{Rep}_k\langle \mathcal{G} \rangle$ are matrix rings over the fixed field $k^{\mathcal{G}}$ and thus*

$$K_*\mathrm{Rep}_k\langle \mathcal{G} \rangle \cong (K_0\mathrm{Rep}_k\langle \mathcal{G} \rangle) \otimes K_*(k^{\mathcal{G}}).$$

In the sequel, define $\mathcal{L} = k^{\mathcal{G}}$.

Proposition 4.5. *There is an isomorphism of rings, $K_0\mathrm{Rep}_k\langle \mathcal{G} \rangle \cong \mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell]$.*

Finally, we compute the homotopy groups of $(K\text{Rep}_k(\mathcal{G}))_\ell^\wedge$ and find that they are isomorphic to those of $(K\mathcal{F})_\ell^\wedge$.

Proposition 4.1. *The homotopy groups of $(\text{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge$ are*

$$\begin{aligned} \pi_n((K\text{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge) &\cong \pi_n((K\mathbb{F}_p)_\ell^\wedge) \oplus \pi_n \Sigma((K\mathbb{F}_p)_\ell^\wedge) \\ &\cong \pi_n((K\mathcal{F})_\ell^\wedge). \end{aligned}$$

In Section 5, we show that the map $\mathcal{E}$ induces the isomorphism of homotopy groups by factoring $\mathcal{E}$ through the intermediate category

$$\text{colim}_{\mathcal{G}'_i \in \text{OpNm}(\mathcal{G})} \text{Rep}_{\mathcal{O}_i}(\mathcal{G}/\mathcal{G}'_i),$$

where $\text{OpNm}(\mathcal{G})$ is defined to be the opposite category of the category of open normal subgroups of $\mathcal{G}$, $\mathcal{O}_i$ is the ring of integers in the fixed field $\mathcal{F}^{\mathcal{G}'_i}$, and $\text{Rep}_{\mathcal{O}_i}(\mathcal{G}/\mathcal{G}'_i)$ is the category of finitely generated $\mathcal{O}_i$ -modules equipped with the continuous semi-linear action by $\mathcal{G}/\mathcal{G}'_i$ on $\mathcal{F}^{\mathcal{G}'_i}$.

Acknowledgments. I would like to thank Tyler Lawson for his many helpful suggestions on a draft of this paper.

2. COMPLETIONS

2.1. The Bousfield-Kan Completion. We briefly review the Bousfield-Kan completion [BK72]. Let $\mathcal{C}$ be a category.

Definition 2.1. A monad (or triple) on $\mathcal{C}$ is a functor $T: \mathcal{C} \rightarrow \mathcal{C}$ together with natural transformations $\eta: \text{id} \rightarrow T$ and $\mu: T \circ T \rightarrow T$ such that for any object $X \in \mathcal{C}$ the diagrams below commute.

$$\begin{array}{ccc} TX & \xrightarrow{T\eta(X)} & T \circ TX & \xleftarrow{\eta(TX)} & TX \\ & \searrow & \downarrow \mu(X) & & \swarrow \\ & & TX & & \end{array} \quad \begin{array}{ccc} T^3(X) & \xrightarrow{\mu(TX)} & T^2(X) \\ T\mu(X) \downarrow & & \downarrow \mu(X) \\ T^2(X) & \xrightarrow{\mu(X)} & T(X) \end{array}$$

See [May72] for more details.

Definition 2.2. Suppose that $c \in \mathcal{C}$. The cosimplicial resolution of c relative to T is the cosimplicial object $\mathcal{R}^\bullet(T, c)$ whose n simplices are obtained by making $n+1$ applications of T . That is, for any nonzero integer n ,

$$\mathcal{R}^n(T, c) := T^{n+1}(c).$$

The coface and codegeneracies are determined by η and μ respectively:

$$\delta^s: T^{k+1}(c) \rightarrow T^{k+2}(c) = T^s(\eta(T^{k+1-s}(c)))$$

and

$$\sigma^t: T^{k+1}(c) \rightarrow T^k(c) = T^t(\mu(T^{k-t-1}(c))).$$

Now let $\mathcal{C}$ be the category of simplicial sets and R a ring. Define F_R to be the composition of the forgetful functor with the free R -module functor applied levelwise. Then F_R (together with the obvious natural transformations) is a monad on $\mathcal{C}$.

Definition 2.3 (Bousfield-Kan). Let $X_\bullet$ be a simplicial set. Then the R -completion of $X_\bullet$, denoted $R_\infty X$, is defined to be the cosimplicial resolution of $X_\bullet$ relative to F_R . The ℓ -adic completion of $X_\bullet$, denoted $X_\ell^\wedge$, is the R -completion for $R = \mathbb{F}_\ell$.

This construction can be extended to the category of spectra.

Definition 2.4. Suppose that $X = \{X_i\}$ is a spectrum. Then the R -completion of X , denoted $R_\infty X$ is defined to be the “spectrification” of the prespectrum $\{R_\infty(X_i)\}$. As before, the $\mathbb{F}_\ell$ -completion is called the ℓ -adic completion and is denoted $X_\ell^\wedge$.

The following propositions are about the homotopy groups of ℓ -adically completed spectra.

Proposition 2.5. *Let X be a spectrum whose homotopy groups are all finitely generated. Then*

$$\pi_n(X_\ell^\wedge) \cong \mathbb{Z}_\ell \otimes \pi_n X.$$

Proof. This follows from Example 5.2 in [BK72]. $\square$

Proposition 2.6. *Let F be a finite field of characteristic p and $L \supseteq F$ an extension of degree prime to ℓ . The inclusion $F \hookrightarrow L$ induces a weak equivalence,*

$$(KF)_\ell^\wedge \xrightarrow{\cong} (KL)_\ell^\wedge.$$

Proof. Let L be a finite extension of F of degree prime to ℓ . The composition of extension and restriction of scalars,

$$KF \longrightarrow KL \longrightarrow KF$$

induces multiplication by $[L : F]$ on $\pi_*(KF)$. Since $[L : F]$ is prime to ℓ , the ℓ -primary components of $\pi_* KF$ are direct summands of the homotopy groups $\pi_* KL$.

Because $\pi_n KF$ and $\pi_n KL$ are both cyclic, they must be isomorphic. $\square$

Proposition 2.7. *The ℓ -adic K -groups of $\mathcal{F}$ are*

$$\pi_*(K\mathcal{F})_\ell^\wedge \cong \pi_*((K\mathbb{F}_p)_\ell^\wedge) \oplus \pi_*\Sigma((K\mathbb{F}_p)_\ell^\wedge).$$

Proof. This is proven in Section 4.2 of [Car07b]. $\square$

2.2. The Derived Completion. To define the derived completion, we observe that when $\gamma: A \rightarrow B$ is a map of commutative ring spectra, then the functor $(B \wedge_A -)$ together with the natural transformations

$$\eta: \text{id} \rightarrow (B \wedge_A -),$$

defined by $\eta = f \wedge \text{id}$ and

$$\mu: (B \wedge_A B \wedge_A -) \rightarrow (B \wedge_A -),$$

defined by multiplication on B , is a monad on the category $\text{Mod}(A)$ of A -modules. When A is the sphere spectrum and B is the Eilenberg-MacLane spectrum of a ring R , the resulting monad $(H(R) \wedge_{\mathbb{S}} -)$ is equivalent in the category of spectra to the monad F_R . Thus for an $\mathbb{S}$ -module M there is a weak equivalence of $\mathbb{S}$ -modules,

$$M_{HR}^\wedge \simeq R_\infty M.$$

Denote by $\tilde{B}$ the functorial fibrant replacement for B in the category of commutative A -algebras, and by $\hat{\mathcal{R}}^\bullet((\tilde{B} \wedge_A -), M)$ the functorial fibrant replacement for $\mathcal{R}^\bullet((\tilde{B} \wedge_A -), M)$ in the category of cosimplicial A -modules.

Definition 2.8. The derived completion $M_\gamma^\wedge$ of M with respect to the map $\gamma: A \rightarrow B$ is the total space,

$$M_\gamma^\wedge := \text{Tot}(\hat{\mathcal{R}}^\bullet(\tilde{B} \wedge_A -, M)).$$

The following propositions and theorems from [Car07a] are the most relevant for our purposes. We refer the reader to [Car07a] for more details.

Proposition 2.9. *Let $A \rightarrow B$ be a map of commutative ring spectra and M an A -module. Then the following statements all hold.*

- (1) *The construction $M \rightarrow M_B^\wedge$ is functorial for homomorphisms of A -modules. The map on B -completions induced by a homomorphism $f: M \rightarrow N$ of A -modules will be denoted by $f_B^\wedge$.*
- (2) *Let $f: M \rightarrow N$ be a homomorphism of A -modules, where A is a commutative ring spectrum, and suppose that $\text{id}_B \wedge_A f$ is a weak equivalence of B -module spectra. Then the map on completions $f_B^\wedge$ is also a weak equivalence of spectra.*
- (3) *Let $M'' \rightarrow M \rightarrow M'$ be a cofibration sequence of A -module spectra. Then the sequence $(M'')_B^\wedge \rightarrow (M)_B^\wedge \rightarrow (M')_B^\wedge$ is a cofibration sequence up to homotopy.*
- (4) *Let $A'' \rightarrow A \rightarrow B$ be a diagram of commutative ring spectra, and let M denote a A -module spectrum. Denote by $M_{A''}$ the module M regarded as an A'' -module. Suppose that the natural map $B \wedge_{A''} (M_{A''}) \rightarrow B \wedge_A M$ is a weak equivalence of spectra. Then the natural map $(M_{A''})_B^\wedge \rightarrow M_B^\wedge$ is also an equivalence of spectra.*
- (5) *There is a natural map $\eta: M \rightarrow M_B^\wedge$.*
- (6) *Suppose M is an A -module spectrum for which the A -module structure admits a B -module structure extending the given A -module structure. Then the natural map $\eta: M \rightarrow M_B^\wedge$ is an equivalence of spectra.*

Theorem 2.10. *Suppose that we have a diagram $A \rightarrow B \xrightarrow{\beta} B'$ of commutative ring spectra and a left A -module spectrum M . Suppose further that A , B , and M are all (-1) -connected, that the homomorphism $\pi_0(\beta)$ is an isomorphism, and that the natural homomorphisms $\pi_0(A) \rightarrow \pi_0(B)$ and $\pi_0(A) \rightarrow \pi_0(B')$ are surjections. Then the natural homomorphism $M_B^\wedge \rightarrow M_{B'}^\wedge$ is an equivalence of A -module spectra.*

Theorem 2.11 (Algebraic to geometric spectral sequence). *Let $A \rightarrow B$ be a map of commutative ring spectra, with $\pi_0 A \rightarrow \pi_0 B$ surjective, and let M be an A -module spectrum. The ring $\pi_0 A$ is commutative, the ring $\pi_0 B$ is a commutative $\pi_0 A$ -algebra, and for each i , the abelian group $\pi_i M$ has a $\pi_0 A$ -module structure. We may therefore construct the derived completion $(\pi_i M)_{\pi_0 B}^\wedge$ for each i . There is a second quadrant spectral sequence with $E_1^{s,t} = \pi_{s+2t}((\pi_{-t} M)_{\pi_0 B}^\wedge)$, converging to $\pi_{t+s}(M_B^\wedge)$.*

The following example from [Car07a] will play an important role in our computation of the homotopy groups $\pi_*((K\text{Rep}_K(\mathcal{G}))_{\alpha_\ell}^\wedge)$.

Example 2.12. Denote by $R[\mathbb{Z}_\ell]$ the ring of isomorphism classes of complex representations of the group of ℓ -adic integers. Then $R[\mathbb{Z}_\ell]$ can be expressed as a colimit

$$R[\mathbb{Z}_\ell] = \operatorname{colim}_n R[\mathbb{Z}/\ell^n\mathbb{Z}].$$

For each finite representation ring, there is an isomorphism,

$$R[\mathbb{Z}/\ell^n\mathbb{Z}] \cong \mathbb{Z}[\mu_{\ell^n}],$$

and thus $R[\mathbb{Z}_\ell] \cong \mathbb{Z}[\mu_{\ell^\infty}]$, where μ_{ℓ^∞} is the group of all ℓ -power roots of unity.

Define α_ℓ as in (1). Then the homotopy groups of the derived completion of the Eilenberg-MacLane spectrum $(HR[\mathbb{Z}_\ell])_{\alpha_\ell}^\wedge$ relative to α_ℓ are

$$\pi_*((HR[\mathbb{Z}_\ell])_{\alpha_\ell}^\wedge) \cong \begin{cases} \mathbb{Z}_\ell & \text{for } i = 0 \text{ or } 1, \\ 0 & \text{otherwise.} \end{cases}$$

3. SEMISIMPLE SKEW GROUP RINGS

In this section we summarize the relevant results of “Semisimple skew group rings and their modules” [LR07], which presents a means of describing the objects, endomorphisms, and tensor products in the category of finitely generated modules over a skew group ring. We will compute the ring K_0 of isomorphism classes of finitely generated modules over a skew group ring under the operations $\oplus$ and $\otimes_E$.

Suppose that E is a field equipped with the action of a finite group G . Then we define the skew group ring $E\langle G \rangle$ to be the E -vector space with basis set G endowed with the multiplication structure determined by $(\alpha g)(\beta h) = \alpha {}^g\beta gh$, for α and $\beta \in E$, for g and $h \in G$, and where ${}^g\beta$ is the image of β under g . If G acts trivially on E , then $E\langle G \rangle$ is the ordinary group ring $E[G]$. In our motivating example, the action by G on E is not trivial; however it is typically not faithful either, and hence the elements of G that act trivially on E form a proper normal subgroup N .

Throughout this section, we will assume that N is abelian and therefore that the action by G on N factors through the quotient G/N , which we will denote by Δ . The assumption that N is abelian also implies that there is a cohomology group $H^2(G/N, N)$. The group G is an extension of G/N by N and thus determines an equivalence class in $H^2(G/N, N)$.

We will also assume that the order of G is invertible in E and thus, by a generalization of Maschke’s Theorem, that $E\langle G \rangle$ is semisimple. Every $E\langle G \rangle$ module can therefore be broken down as a direct sum of simple modules. Every simple module is isomorphic to a simple module over a single factor in the Artin-Wedderburn decomposition of $E\langle G \rangle$ as a product of matrix rings

$$E\langle G \rangle = \prod_{i=1}^s M(d_i, D_i),$$

where s and d_i are positive integers and D_i is a division ring (which will contain the fixed field $F := E^G$). This decomposition is unique; therefore, describing the objects and endomorphisms of $E\langle G \rangle$ -modules is equivalent to finding the division rings D_i and the corresponding dimensions d_i . Each D_i is a central simple algebra over its center Z_i , so D_i corresponds to an element of $\operatorname{Br}(Z_i)$.

The product $Z = \prod Z_i$ is the center of $E\langle G \rangle$.

Proposition 3.1. *The map*

$$\mu: E \otimes_F Z \longrightarrow E[N]$$

defined by

$$e \otimes z \mapsto ez,$$

where $e \in E$ and $z \in Z$, is an E -algebra isomorphism.

Now define $\text{Br}(Z) := \prod \text{Br}(Z_i)$. By a theorem of Auslander and Goldman [AG60] we can identify $\text{Br}(Z)$ with the cohomology group $H^2(G_F, (\overline{F} \otimes Z)^*)$. The skew group ring $E\langle G \rangle$ thus corresponds to a 2-cocycle in $H^2(G_F, (\overline{F} \otimes Z)^*)$. There is a natural map obtained by composing the maps

$$\begin{aligned} H^2(G/N, N) &\xrightarrow{\cong} H^2(\text{Gal}(E/F), N) \\ &\xrightarrow{\text{inf.}} H^2(G_F, N) \\ &\xrightarrow{\text{inc.}} H^2(G_F, F[N]^*) \\ &\xrightarrow{\mu} H^2(G_F, (\overline{F} \otimes Z)^*) \\ &\xrightarrow{\cong} \text{Br}(Z), \end{aligned}$$

where the map “inf.” is inflation and “inc.” is inclusion. Call this composite Φ .

Theorem 3.2. *The map Φ sends $[G] \in H^2(G/N, N)$ to $[E\langle G \rangle] \in \text{Br}(Z)$.*

Corollary 3.3. *If $\text{Br}(F) = 1$, there is an isomorphism of rings,*

$$E\langle G \rangle \cong M([E : F], Z).$$

Proposition 3.4. *As a Z -module, $E\langle G \rangle$ is free of rank $[E : F]^2$.*

This describes $E\langle G \rangle$ up to isomorphism.

To compute the multiplicative structure of $K_0(E\langle G \rangle)$, we first observe that the set $\text{Hom}_F(Z, \overline{F})$ of F -algebra homomorphisms is endowed with a natural action by the absolute Galois group G_F which is defined by

$$(g * \phi)(z) := g(\phi(z))$$

for $g \in G_F$ and $\phi \in \text{Hom}_F(Z, \overline{F})$. This action partitions $\text{Hom}_F(Z, \overline{F})$ into a disjoint union of orbits $\text{Hom}_F(Z_i, \overline{F}) = \coprod \text{Hom}(Z_i, \overline{F})$.

Corollary 3.5. *The map μ induces a bijection of sets*

$$\mu_*: \text{Hom}(Z, \overline{F}) \xrightarrow{\cong} \text{Hom}(N, \overline{F}^*)$$

that is G_F equivariant if the action by G_F on $\text{Hom}(N, \overline{F}^*)$ is defined in the natural way,

$$(3) \quad (g * \phi)(n) := g(\phi(\overline{g}^{-1}n)),$$

where $g \in G_F$, $\phi \in \text{Hom}(N, \overline{F}^*)$, and $n \in N$; the element $\overline{g}$ is the image of g in the quotient $G/N \cong \text{Gal}(E/F)$ of G_F and $\overline{g}^{-1}n$ is the image of n under conjugation by $\overline{g}^{-1}$.

Under μ_* , the orbits $\text{Hom}(Z_i, \overline{F})$ get sent to orbits of $\text{Hom}(N, \overline{F}^*)$ which we call $\mathfrak{D}_i$; each of the $\mathfrak{D}_i$ therefore corresponds to a factor $M(d_i, D_i)$ of $E\langle G \rangle$. Note that the field Z_i is isomorphic to a fixed field of $\overline{F}$ under the subgroup $\text{Stab}_{G_F}(\chi)$ of elements in G_F that fix any character χ in $\mathfrak{D}_i$. The degree of Z_i over F is equal to the number of elements in $\mathfrak{D}_i$.

Now fix a common separable closure $\overline{F}$ of F and all separable extensions of F , and write $N^\vee := \text{Hom}(N, \overline{F}^*)$. We will associate to every $E\langle G \rangle$ -module W an element of the group ring $\mathbb{Z}[N^\vee]$ as follows.

We start by restricting W to an E -linear representation of N . Since N is abelian, we can write $\overline{F} \otimes_E W$ as a sum of 1-dimensional representations

$$\overline{F} \otimes_E W \cong \bigoplus_{N^\vee} U_\chi^{m_\chi},$$

where $m_\chi \in \mathbb{Z}^{\geq 0}$. This decomposition is unique and suggests a map ρ defined by

$$(4) \quad \rho(W) := \sum m_\chi \chi$$

from $K_0 E\langle G \rangle$ to the group ring $\mathbb{Z}[N^\vee]$.

Proposition 3.6. *For every $E\langle G \rangle$ -module W , the image $\rho(W)$ is stable under the action by G_F defined in equation (3).*

Proposition 3.7. *For a simple module V_i , the set of elements that appear with a nonzero coefficient in $\rho(V_i)$ is exactly the orbit $\mathfrak{D}_i$ of $N^\vee$, defined in Corollary 3.5. The map ρ is therefore injective.*

If V_i is a simple module over the i th factor $M(d_i, D_i)$ of $E\langle G \rangle$, then the center Z_j of a different factor, where $1 \leq j \leq s$ and $j \neq i$, acts trivially on V_i . The elements of Z_i , on the other hand, act through the characters in $\text{Hom}(Z_i, \overline{F})$ by

$$z \cdot v = \chi(z)v,$$

where $z \in Z_i$, $v \in V$, and $\chi \in \text{Hom}(Z_i, \overline{F})$. By Corollary 3.5, these characters correspond exactly to those in $\mathfrak{D}_i$.

It follows immediately from Propositions 3.6 and 3.7 that for a simple module V_i , there is some positive integer m_i , which we call the “multiplicity” of V_i , such that

$$\rho(V_i) = m_i \sum_{\chi \in \mathfrak{D}_i} \chi.$$

Corollary 3.8. *If $\text{Br}(F) = 1$, then $m_i = 1$ for all irreducible $E\langle G \rangle$ -modules V_i .*

Theorem 3.9. *The map ρ is multiplicative with respect to $\otimes_E$, and hence an $E\langle G \rangle$ -module W is irreducible exactly when its image in $\mathbb{Z}[N^\vee]$ is*

$$\rho(W) = m_i \sum_{\chi \in \mathfrak{D}_i} \chi,$$

for some i . If W' is another module over $E\langle G \rangle$, then

$$W \otimes_E W' \cong \bigoplus_{\mathcal{V}} V_i^{r_i}$$

where $r_i \in \mathbb{Z}^{\geq 0}$ and $\mathcal{V}$ is a set containing one representative V_i for each isomorphism class of simple modules, exactly when $\rho(W) \cdot \rho(W') = \sum_{\mathcal{V}} \rho(V_i^{r_i})$.

Example 3.10. Suppose that the Brauer group $\text{Br}(F)$ is trivial and therefore that $\text{Hom}(N, \overline{F}^*)$ breaks into singleton orbits under the action by G_F . Then the map ρ induces an isomorphism of rings

$$\rho : K_0(E\langle G \rangle) \xrightarrow{\cong} \mathbb{Z}[N^\vee].$$

Example 3.11. We will show in Proposition 4.1 that $\mathcal{G} \cong \mathbb{Z}_\ell \rtimes_\xi \mathbb{Z}_\ell$ where the action is the Frobenius action. Let $\mathcal{G}_i = \mathbb{Z}/\ell^i\mathbb{Z} \rtimes \mathbb{Z}/\ell^{i'}\mathbb{Z}$ be the quotient of the group $\mathcal{G}$ by the subgroup $\mathcal{G}'_i = \ell^i\mathbb{Z}_\ell \rtimes \ell^{i'}\mathbb{Z}_\ell$, where $i > i'$ are positive integers. The integer i' is defined to be the degree of the Galois extension $\mathcal{L}(\zeta)/\mathcal{L}$, where ζ is a primitive $\ell^{i'}$ th root of unity. The group $\mathcal{G}_i$ acts on $(\overline{\mathbb{F}}_p)^{\ell^i\mathbb{Z}_\ell}$ (which we denote k_i) in the obvious way, and the fixed field of this action is $\mathcal{L}$.

Let $\mathcal{N}_i := \mathbb{Z}/\ell^i\mathbb{Z}$ be the subgroup of elements of $\mathcal{G}_i$ that act trivially on k_i and define Δ_i to be the quotient $\mathbb{Z}/\ell^{i'}\mathbb{Z} \cong \mathcal{G}_i/\mathcal{N}_i$. Let η be a generator of $\mathcal{N}_i$. The quotient group Δ_i (which is also a subgroup of $\mathcal{G}$) is generated by the Frobenius automorphism ϕ and sends an automorphism $\eta^r \in \mathcal{N}_i$, where r is an integer between 0 and $\ell^{i'} - 1$, to its p th power; that is, $\phi\eta^r\phi^{-1} = \eta^{pr}$. The action by the absolute Galois group $G_{\mathcal{L}}$ on $\mathcal{N}_i^\vee = \text{Hom}(\mathcal{N}_i, \overline{\mathcal{L}}^*)$ divides $\mathcal{N}_i^\vee$ into singleton orbits. By Example 3.10,

$$K_0\left(k_i\langle \mathbb{Z}/\ell^i\mathbb{Z} \rtimes \mathbb{Z}/\ell^{i'}\mathbb{Z} \rangle\right) \cong \mathbb{Z}[\mu_{\ell^i}].$$

Now we consider the case where G is a profinite group. Define $\text{OpNm}(G)$ to be the category whose objects are the open normal subgroups of G . For every pair G'' and G' of such subgroups, the morphism set $\text{Mor}(G'', G')$ has exactly one element if G' is a subgroup of G'' and is empty otherwise. Let $\text{OpNm}(G)^{\text{op}}$ be the opposite category. The functor

$$\text{Rep}_E\langle G/- \rangle : \text{OpNm}(G)^{\text{op}} \rightarrow \text{ExactCats}$$

takes an open normal subgroup $G' \subseteq G$ to the exact category $\text{Rep}_{E'}\langle G/G' \rangle$, where $E' = E^{G'}$.

Proposition 3.12. *The category $\text{Rep}_E\langle G \rangle$ is a colimit*

$$\text{Rep}_E\langle G \rangle \cong \text{colim}_{G' \in \text{OpNm}(G)^{\text{op}}} \text{Rep}_{E'}\langle G/G' \rangle.$$

Each constituent $\text{Rep}_{E'}\langle G/G' \rangle$ of the colimit may be identified with a subcategory of $\text{Rep}_E\langle G \rangle$ under the functor

$$\varphi' : \text{Rep}_{E'}\langle G/G' \rangle \rightarrow \text{Rep}_E\langle G \rangle,$$

which is defined by

$$\varphi'(W) = E \otimes_{E'} W.$$

The action by an element $g \in G$ on an element $e \otimes w \in E \otimes W$ is defined by

$$g(e \otimes w) = {}^g e \otimes \overline{g}(w),$$

where ${}^g e$ is the image of e under the action by g , the element $\overline{g}$ is the image of g under the quotient map $G \rightarrow G/G'$, and $\overline{g}(w)$ is the image of w under the action by $\overline{g}$.

Corollary 3.13. *There is an isomorphism of rings,*

$$K_0\text{Rep}_E\langle G \rangle \cong \text{colim}_{G' \in \text{OpNm}(G)} K_0E'\langle G/G' \rangle.$$

4. THE HOMOTOPY GROUPS OF $(K\text{Rep}_k\langle\mathcal{G}\rangle)_{\alpha_\ell}^\wedge$

In this section we compute the homotopy groups of $K\text{Rep}_k\langle\mathcal{G}\rangle$, which we will feed in to the algebraic to geometric spectral sequence to obtain the homotopy groups of the derived completion $(K\text{Rep}_k\langle\mathcal{G}\rangle)_{\alpha_\ell}^\wedge$.

Define $\mathbb{N}$ to be the filtered category whose objects are the natural numbers and whose morphism sets $\text{Mor}_{\mathbb{N}}(m, n)$ consist of one element when n divides m and are empty otherwise. For any ring R and any positive integer r , we will denote by $R[[x^{(1/r)}, x^{(1/s)}]]$ the colimit,

$$R[[x^{(1/r)}, x^{(1/s)}]] := \text{colim}_{i \in \mathbb{N} \text{ s.t. } r, s \nmid i} R[[x^{1/i}]].$$

If R is already a colimit of the form

$$R = \text{colim}_{i \in \mathcal{J}} R'[[x^{1/i}]]$$

for some ring R' and some subset $\mathcal{J} \subseteq \mathbb{N}$, then we denote by $R[x^{(1/r)}]$ the colimit

$$R[x^{(1/r)}] := \text{colim}_{i \in \mathcal{J} \text{ s.t. } r \nmid i} R'[[x^{1/i}]].$$

We define $F((x^{(1/r)}))$ and $F(x^{(1/r)})$ in the analogous way.

We start by describing the tame Galois group $\mathcal{G}$. It is easy to see that there is a normal subgroup of $\mathcal{G}$, which we denote by $\mathcal{N}$, that is isomorphic to $\text{Gal}(\overline{\mathcal{L}}/\mathcal{L}) \cong \mathbb{Z}_\ell$. The quotient by this normal subgroup, which permutes the roots of x , will be denoted Δ .

Proposition 4.1. *The tame Galois group of $\mathcal{F}$ is the semidirect product $\mathbb{Z}_\ell \rtimes \mathbb{Z}_\ell$.*

Proof. The quotient Δ sits inside $\mathcal{G}$ as the Galois group $\Delta \cong \text{Gal}(\mathcal{F}^t/\mathcal{F}^t((x^{(1/\ell)})))$. The subgroups $\mathcal{N}$ and Δ intersect trivially and their product fixes exactly $\mathcal{F}$. Therefore $\mathcal{G}$ is the product of subgroups $\mathcal{N} \cdot \Delta$. $\square$

Remark 4.2. The generator of Δ (the Frobenius automorphism) acts on $\mathcal{N}$ by raising elements to the ℓ -th power.

Remark 4.3. The subgroup $\mathbb{Z}_\ell \rtimes \mathbb{Z}_\ell$ is not necessarily normal in $G_{\mathbb{F}_p((x))}^t$. This is because for all primes q different from ℓ and p , $\mathcal{F}$ contains all q th roots of x but not necessarily all such roots of 1. For example, if $q \mid p^\ell - 1$, then the order of p modulo q will be ℓ and hence the q th roots of unity can only be adjoined by making a degree ℓ extension. The extension $\mathcal{F}/\mathbb{F}_p((x))$ is therefore not necessarily Galois.

Define $\mathcal{G}_i$, $\mathcal{G}'_i$, $\mathcal{N}_i$, and k_i as in Example 3.11. By Proposition 3.12 there is an equivalence of categories

$$\text{Rep}_k\langle\mathcal{G}\rangle \cong \text{colim}_{\mathcal{G}'_i \in \text{OpNm}(\mathcal{G})} \text{Rep}_{k_i}\langle\mathcal{G}/\mathcal{G}'_i\rangle.$$

Proposition 4.4. *The ring of endomorphisms of every irreducible object in $\text{Rep}_k\langle\mathcal{G}\rangle$ is $\mathcal{L}$. The K -groups of $\text{Rep}_k\langle\mathcal{G}\rangle$ are therefore*

$$K_*\text{Rep}_k\langle\mathcal{G}\rangle \cong (K_0\text{Rep}_k\langle\mathcal{G}\rangle) \otimes K_*\mathcal{L}.$$

Proof. The orbits of $\text{Hom}(\mathcal{N}_i, k^*)$ under the action by the absolute Galois group $G_{\mathcal{L}}$ are all singleton orbits, and thus by Proposition 3.7, there is a one-to-one correspondence between characters of $\mathcal{N}_i$ and irreducible representations of $\mathcal{G}'_i$. An

irreducible representation V that corresponds to the character $\chi \in \text{Hom}(\mathcal{N}_i, \overline{k_i}^*)$ is a k_i -vector space on which $\mathcal{N}_i$ acts by

$$n \cdot v = \chi(n)v,$$

for $n \in \mathcal{N}_i$ and $v \in V$. Since $\mathcal{G}_i$ is a semidirect product of $\mathcal{N}_i$ and Δ_i , every $g \in \mathcal{G}'_i$ can be written uniquely as a product $g = nd$, for $n \in \mathcal{N}_i$ and $d \in \Delta_i$. The action by g on $ev \in V$ is determined by

$$g(ev) = {}^d e \cdot \chi(n) \cdot d(v).$$

The endomorphisms of V as a k_i -vector space are just k_i ; the endomorphisms of V that commute with $\mathcal{G}_i$ are exactly those that are fixed by the action of Δ_i , namely, the elements of $\mathcal{L}$.

Now since every object in $\text{Rep}_k(\mathcal{G})$ has a canonical decomposition as a direct sum of irreducible representations, the result follows. $\square$

Proposition 4.5. *The ring $K_0\text{Rep}_k(\mathcal{G})$ of isomorphism classes of k -semilinear representations of $\mathcal{G}$ is*

$$K_0\text{Rep}_k(\mathcal{G}) \cong \mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell].$$

Proof. This follows immediately from Example 3.11 and Proposition 3.12. $\square$

Now that we have computed the homotopy groups of $K\text{Rep}_k(\mathcal{G})$, we will use them to compute, in Proposition 4.9, the homotopy groups of the derived completion $(K\text{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge$. We will use the following well-known lemma which describes the homotopy groups of the ℓ -adic completion $(K\mathcal{L})_\ell^\wedge$.

Lemma 4.6. *Let X be a spectrum with homotopy groups $M_n := \pi_n X$, and let $U_n \subseteq M_n$ be a uniquely ℓ -divisible part of M_n . Then if M_n/U_n is a finitely generated abelian group, the homotopy groups of the ℓ -adic completion of X are*

$$\pi_n(X_\ell^\wedge) \cong M_n/U_n \otimes \mathbb{Z}_\ell.$$

The next two technical lemmas simplify our computations of the homotopy groups of derived completions.

Lemma 4.7. *Let $\alpha: \mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell] \rightarrow \mathbb{F}_\ell$ be the mod- ℓ augmentation map and let X be an $H(\mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell])$ -module. Let $M_n := \pi_n(X)$, and define $U_n \subseteq M_n$ to be the uniquely ℓ -divisible part of M_n . Then if M_n/U_n is finitely generated, the map $\beta: X \rightarrow X_\ell^\wedge$ induces a weak equivalence*

$$(\mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell] \otimes \pi_* X)_\alpha^\wedge \xrightarrow{\cong} \left(\mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell] \otimes \pi_*(X_\ell^\wedge) \right)_\alpha^\wedge$$

on derived completions.

Proof. By Lemma 4.6,

$$\pi_*(X_\ell^\wedge) \cong M_n/U_n \otimes \mathbb{Z}_\ell.$$

There is an isomorphism

$$\mathbb{F}_\ell \otimes M_n \cong \mathbb{F}_\ell \otimes M_n/U_n \otimes \mathbb{Z}_\ell.$$

and hence, by a theorem of Elmendorf-Kriz-Mandell-May [EKMM97], β induces weak equivalences between the n -simplices of the derived completions. $\square$

Lemma 4.8. *Let $R \rightarrow S$ be a map of commutative rings and M an R -module with no $\mathbb{Z}$ -torsion. Then if A is a finitely generated abelian group,*

$$\pi_* \left((M \otimes A)_S^\wedge \right) \cong \pi_*(M_S^\wedge) \otimes A.$$

Proof. Let $0 \rightarrow \mathbb{Z}^n \xrightarrow{i} \mathbb{Z}^m \rightarrow A \rightarrow 0$ be a free resolution of A . The result now follows immediately from the fact that the derived completion and π_* preserve fiber sequences. $\square$

Proposition 4.9. *The algebraic to geometric spectral sequence for $(K\text{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge$ collapses at the E_1 -term and thus $(K\text{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge$ has homotopy groups*

$$\pi_i \left((K\text{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge \right) \cong \pi_i \left((K\mathbb{F}_p)_\ell^\wedge \right) \oplus \pi_{i-1} \left((K\mathbb{F}_p)_\ell^\wedge \right).$$

Proof. The E_1 -term of the algebraic to geometric spectral sequence is

$$E_1^{s,t} = \pi_{s+2t} \left(\left(\mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell] \otimes \pi_{-t} K\mathcal{L} \right)_{\mathbb{F}_\ell}^\wedge \right).$$

By Lemmas 4.6 and 4.8 along with Example 2.12, this becomes

$$E_1^{s,t} = \begin{cases} \pi_{-t} \left((K\mathcal{L})_\ell^\wedge \right) & \text{when } s + 2t = 0 \text{ or } 1 \\ 0 & \text{otherwise.} \end{cases}$$

Our E_1 page is therefore as in the diagram below.

The diagram shows the E_1 page of the spectral sequence. The horizontal axis is labeled t and the vertical axis is labeled s . The terms are arranged in a grid. The diagonal terms are labeled with the derived completion of the tensor product of the group ring and the K-theory of the field. The off-diagonal terms are zero. The diagonal terms are: at $(0,0)$ is $\pi_0((K\mathcal{L})_\ell^\wedge)$; at $(1,0)$ is $\pi_1((K\mathcal{L})_\ell^\wedge)$; at $(2,0)$ is $\pi_1((K\mathcal{L})_\ell^\wedge)$; at $(3,0)$ is $\pi_2((K\mathcal{L})_\ell^\wedge)$; at $(4,0)$ is $\pi_2((K\mathcal{L})_\ell^\wedge)$; at $(0,1)$ is $\pi_0((K\mathcal{L})_\ell^\wedge)$; at $(1,1)$ is $\pi_1((K\mathcal{L})_\ell^\wedge)$; at $(2,1)$ is $\pi_1((K\mathcal{L})_\ell^\wedge)$; at $(3,1)$ is $\pi_2((K\mathcal{L})_\ell^\wedge)$; at $(4,1)$ is $\pi_2((K\mathcal{L})_\ell^\wedge)$. All other terms are 0.

By Proposition 2.6, the diagonal lines each correspond to a copy of $\pi_*((K\mathbb{F}_p)_\ell)^\wedge$. All of the differentials vanish, and hence the spectral sequence collapses. Therefore the homotopy groups of the derived completion are

$$\pi_i\left((K\mathrm{Rep}_k\langle\mathcal{G}\rangle)_{\alpha_\ell}^\wedge\right) \cong \pi_i\left((K\mathbb{F}_p)_\ell^\wedge\right) \oplus \pi_{i-1}\left((K\mathbb{F}_p)_\ell^\wedge\right).$$

□

5. THE MAP $\mathcal{E}$

In the proof of Theorem 5.4, we will follow a strategy similar to the one found in Section 4 of [Car07b]. Define k_i , $\mathcal{G}'_i$ and $\mathcal{G}_i$ as in example 3.11, let $\mathcal{F}_i := (\mathcal{F}^t)^{\mathcal{G}'_i}$ and let $\mathcal{O}_i$ be the ring of integers in $\mathcal{F}_i$. Denote by $\mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle$ the category of finitely generated $\mathcal{O}_i$ -modules equipped with the continuous semilinear action by $\mathcal{G}_i$ that is inherited from the natural action by $\mathcal{G}_i$ on $\mathcal{F}_i$.

The categories $\mathrm{colim} \mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle$ and $\mathrm{colim} \mathrm{Rep}_{\mathcal{F}_i}\langle\mathcal{G}_i\rangle$ are closed under tensor products with objects in $\mathrm{Rep}_k\langle\mathcal{G}\rangle$, and thus their K -theory spectra are both $K\mathrm{Rep}_k\langle\mathcal{G}\rangle$ -modules. We may therefore take their derived completions across the augmentation map $\alpha_\ell : K\mathrm{Rep}_k\langle\mathcal{G}\rangle \rightarrow K\mathfrak{F}_\ell$ (equation 1). As in [Car07b], we will show that the intermediate category $\mathrm{colim} \mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle$ satisfies the two conditions below. These conditions will be proven in Propositions 5.2 and 5.3, respectively.

- (1) Extension of scalars induces a weak equivalence of completed K -theory spectra,

$$(K\mathrm{Rep}_k\langle\mathcal{G}\rangle)_{\alpha_\ell}^\wedge \xrightarrow{\sim} (\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle)_{\alpha_\ell}^\wedge.$$

- (2) Extension of scalars induces a weak equivalence of completed K -theory spectra,

$$(\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle)_{\alpha_\ell}^\wedge \xrightarrow{\sim} (K\mathrm{Rep}_{\mathcal{F}^t}\langle\mathcal{G}\rangle)_{\alpha_\ell}^\wedge \simeq (K\mathcal{F})_{\alpha_\ell}^\wedge.$$

We start by proving a fact about K -theory spectra of rings of the form $R = E[[x^{\frac{1}{r}}]]\langle G \rangle$ where E is a field, G is a finite group, and r is a positive integer. Denote by $\mathrm{Mod}(R)$ the category of finitely generated projective R -modules, and by $\mathrm{NoTor}(R) \subseteq \mathrm{Mod}(R)$ the full subcategory of finitely generated torsion free projective R -modules.

Lemma 5.1. *The inclusion functor $\mathrm{NoTor}(R) \hookrightarrow \mathrm{Mod}(R)$ induces a weak equivalence of K -theory spectra.*

Proof. Let M be a projective R -module on n generators, where n is a positive integer. Then we can construct a surjective map $R^n \rightarrow M$. The kernel of this map is torsion free, and thus we have a length 2 resolution of M by torsion free modules. The result now follows by “Reduction by Resolution,” Theorem 3.3 of [Qui73]. □

Proposition 5.2. *Extension of scalars induces a weak equivalence,*

$$(\mathrm{colim} K\mathrm{Rep}_{k_i}\langle\mathcal{G}_i\rangle)_{\alpha_\ell}^\wedge \xrightarrow{\sim} (\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle)_{\alpha_\ell}^\wedge.$$

Proof. First we show that the map $t_i : \mathrm{Rep}_{k_i}\langle\mathcal{G}_i\rangle \xrightarrow{\mathcal{O}_i\langle\mathcal{G}_i\rangle \otimes_{k_i\langle\mathcal{G}_i\rangle} (-)} \mathrm{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle$ induces an isomorphism on π_0 .

The categories $\text{Rep}_{k_i}\langle\mathcal{G}_i\rangle$ and $\text{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle$ can be identified with the categories $\text{Mod}(k_i\langle\mathcal{G}_i\rangle)$ and $\text{Mod}(\mathcal{O}_i\langle\mathcal{G}_i\rangle)$, respectively, since the group $\mathcal{G}_i$ is finite. By Lemma 5.1, it suffices to show that t_i induces an isomorphism

$$\tilde{t}_i: \pi_0 K\text{NoTor}(k_i\langle\mathcal{G}_i\rangle) \xrightarrow{\cong} \pi_0 K\text{NoTor}(\mathcal{O}_i\langle\mathcal{G}_i\rangle).$$

Clearly, $\tilde{t}_i$ is an injection on the set of isomorphism classes of objects. To see that it is a surjection, we start with a module M over the ring $\mathcal{O}_i\langle\mathcal{G}_i\rangle$. Let I be the kernel of the quotient map $\mathcal{O}_i\langle\mathcal{G}_i\rangle \rightarrow k_i\langle\mathcal{G}_i\rangle$; I is generated by the fractional powers of x . The module IM is closed under multiplication by $\mathcal{O}_i\langle\mathcal{G}_i\rangle$, and hence the quotient map $M \xrightarrow{\gamma} M/IM$ is a map of $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ -modules. Both M and M/IM are, by restriction, modules over $k_i\langle\mathcal{G}_i\rangle$ as well, and since $k_i\langle\mathcal{G}_i\rangle$ is semisimple, the map γ splits. Therefore we have a section $s: M/IM \rightarrow M$. This map can be extended to a map of $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ -modules,

$$S: \mathcal{O}_i\langle\mathcal{G}_i\rangle \otimes_{k_i\langle\mathcal{G}_i\rangle} M/IM \rightarrow M.$$

Now as a consequence of Nakayama's Lemma, a homomorphism $f: X \rightarrow Y$ of free modules over a complete local ring with maximal ideal M is an isomorphism exactly when the induced map on the quotient is.

Since M and $\mathcal{O}_i\langle\mathcal{G}_i\rangle \otimes_{k_i\langle\mathcal{G}_i\rangle} M/IM$ are free over $\mathcal{O}_i$ and the induced map on quotients

$$(\mathcal{O}_i\langle\mathcal{G}_i\rangle \otimes_{k_i\langle\mathcal{G}_i\rangle} M/IM) / I(\mathcal{O}_i\langle\mathcal{G}_i\rangle \otimes_{k_i\langle\mathcal{G}_i\rangle} M/IM) \rightarrow M/IM$$

is an isomorphism, the map S must be an isomorphism of $\mathcal{O}_i$ -modules. Thus every module over $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ is "extended" from $k_i\langle\mathcal{G}_i\rangle$.

Next we show that the endomorphism rings of the irreducible modules over $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ are of the form $\mathcal{L}[[x^{(1/p)}, x^{(1/\ell^{i+1})}]]$ by examining the irreducible $k_i\langle\mathcal{G}_i\rangle$ -modules.

By Example 3.11 the irreducible $k_i[\mathcal{G}_i]$ -modules are in bijective correspondence with the group $\text{Hom}(\mathcal{N}_i, k)$, where $\mathcal{N}_i = \mathbb{Z}/\ell^i\mathbb{Z}$. Suppose that M_α is an irreducible $k_i\langle\mathcal{G}_i\rangle$ module corresponding to a character $\alpha \in \text{Hom}(\mathcal{N}_i, k)$. Then because $\mathcal{G}_i$ acts on M_α through α , the tensor product $M'_\alpha := \mathcal{O}_i\langle\mathcal{G}_i\rangle \otimes_{k_i\langle\mathcal{G}_i\rangle} M_\alpha$ is isomorphic as an $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ -module to $\mathcal{O}_i$. The endomorphisms of M'_α over $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ are exactly the elements of M'_α that commute with the action by $\mathcal{O}_i\langle\mathcal{G}_i\rangle$ —namely, $\mathcal{L}[[x^{(1/p)}, x^{(1/\ell^{i+1})}]]$.

The homotopy groups of $K\text{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle$ are therefore

$$\pi_* K\text{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle = K_0\text{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle \otimes K_*\mathcal{L}[[x^{(1/p)}, x^{(1/\ell^{i+1})}]],$$

and thus the homotopy groups of the colimit are

$$\pi_* (\text{colim } K\text{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle) = \mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell] \otimes K_*\mathcal{L}[[x^{(1/p)}]].$$

We now have the commutative diagram below. We set $\mathfrak{Z} := \mathbb{Z}[\mathbb{Q}_\ell/\mathbb{Z}_\ell]$ in all of the diagrams of this proof for typesetting purposes.

$$\begin{array}{ccc} \pi_* K\text{Rep}_k\langle\mathcal{G}\rangle \cong \mathfrak{Z} \otimes \pi_* K\mathcal{L} & \xrightarrow{\quad} & \mathfrak{Z} \otimes \pi_* ((K\mathcal{L})_\ell^\wedge) \\ \downarrow t_* & & \downarrow (t_*)_\ell^\wedge \\ \pi_* \text{colim } K\text{Rep}_{\mathcal{O}_i}\langle\mathcal{G}_i\rangle \cong \mathfrak{Z} \otimes \pi_* K\mathcal{L}[[x^{(1/p)}]] & \rightarrow & \mathfrak{Z} \otimes \pi_* ((K\mathcal{L}[[x^{(1/p)}]])_\ell^\wedge) \\ \downarrow q & & \cong \downarrow (q)_\ell^\wedge \\ \pi_* K\text{Rep}_k\langle\mathcal{G}\rangle \cong \mathfrak{Z} \otimes K\mathcal{L} & \xrightarrow{\quad} & \mathfrak{Z} \otimes \pi_* ((K\mathcal{L})_\ell^\wedge) \end{array}$$

The horizontal arrows are induced by the natural maps that come from the Bousfield completion, the map t is defined to be the colimit of the t_i , and the map t_* is the induced map on homotopy groups. The map q is induced by the quotient map $\mathcal{L}[[x^{(1/p)}]] \rightarrow \mathcal{L}$ and, by a theorem of Suslin [Sus84], induces a weak equivalence on the ℓ -adic completions. The composition $q \circ t$ is the identity and hence if $(q)_\ell^\wedge$ is a weak equivalence, $(t_*)_\ell^\wedge$ must also be a weak equivalence. The maps in the top two rows of the diagram above induce the diagram of E_1 -terms for the algebraic to geometric spectral sequence below.

$$\begin{array}{ccc} \pi_{s+2t} \left(\left(\mathfrak{Z} \otimes \pi_{-t} K\mathcal{L} \right)_{\mathbb{F}_\ell}^\wedge \right) & \xrightarrow{\cong} & \pi_{s+2t} \left(\left(\mathfrak{Z} \otimes \pi_{-t} K\mathcal{L}_\ell^\wedge \right)_{\mathbb{F}_\ell}^\wedge \right) \\ \downarrow \varepsilon(t_*) & & \downarrow \varepsilon(t_*)_\ell^\wedge \\ \pi_{s+2t} \left(\left(\mathfrak{Z} \otimes \pi_{-t} K\mathcal{L}[[x^{(1/p)}]] \right)_{\mathbb{F}_\ell}^\wedge \right) & \xrightarrow{\cong} & \pi_{s+2t} \left(\left(\mathfrak{Z} \otimes \pi_{-t} K\mathcal{L}[[x^{(1/p)}]]_\ell^\wedge \right)_{\mathbb{F}_\ell}^\wedge \right). \end{array}$$

By Lemma 4.7 the horizontal maps are isomorphisms on E_1 -terms. Therefore the map $\varepsilon(t_*)$ of E_1 -terms is an isomorphism. Since by Lemma 4.9, the spectral sequence collapses at E_1 , the map t induces a weak equivalence,

$$(K\mathrm{Rep}_k\langle \mathcal{G} \rangle)_{\alpha_\ell}^\wedge \xrightarrow{\cong} (\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle)_{\alpha_\ell}^\wedge.$$

□

Proposition 5.3. *Extension of scalars induces a weak equivalence of completed K -theory spectra,*

$$(\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle)_{\alpha_\ell}^\wedge \xrightarrow{\cong} (K\mathrm{Rep}_{\mathcal{F}^t}\langle \mathcal{G} \rangle)_{\alpha_\ell}^\wedge \cong K\mathcal{F}_{\alpha_\ell}^\wedge.$$

Proof. As previously mentioned, the weak equivalence $(K\mathrm{Rep}_{\mathcal{F}^t}\langle \mathcal{G} \rangle)_{\alpha_\ell}^\wedge \simeq (K\mathcal{F})_{\alpha_\ell}^\wedge$ is an immediate consequence of the equivalence of categories $\mathrm{Rep}_{\mathcal{F}^t}\langle \mathcal{G} \rangle \cong \mathrm{Mod}(\mathcal{F})$. Therefore we need only show that

$$(\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle)_{\alpha_\ell}^\wedge \rightarrow (K\mathrm{Rep}_{\mathcal{F}^t}\langle \mathcal{G} \rangle)_{\alpha_\ell}^\wedge$$

is a weak equivalence. Since $\mathcal{G}_i$ is finite, the category $\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle$ can also be thought of as a category of finitely generated modules over the skew group ring $\mathcal{O}_i\langle \mathcal{G}_i \rangle$. In this ring, inverting all powers of x (including fractional ones) yields the ring $\mathcal{F}_i$, and hence we can use Corollary 5.12 from [Swa68] and Quillen's localization theorem [Qui73] to conclude that there is a homotopy fiber sequence of K -theory spectra,

$$K\mathcal{T}_i \rightarrow K\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle \rightarrow K\mathrm{Rep}_{\mathcal{F}_i}\langle \mathcal{G}_i \rangle,$$

where $\mathcal{T}_i$ is the full subcategory of modules in $\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle$ that localize to 0. These are precisely the modules that are annihilated by some (possibly fractional) power of x .

The category $\mathrm{colim} \mathcal{T}_i$ is closed under the tensor product with objects in $\mathrm{Rep}_k\langle \mathcal{G} \rangle$. We may therefore take its derived completion across the map α_ℓ (see equation 1). Since colimits and derived completions preserve fiber sequences, we have a fiber sequence of $K\mathrm{Rep}_k\langle \mathcal{G} \rangle$ -modules

$$(\mathrm{colim} K\mathcal{T}_i)_{\alpha_\ell}^\wedge \rightarrow (\mathrm{colim} K\mathrm{Rep}_{\mathcal{O}_i}\langle \mathcal{G}_i \rangle)_{\alpha_\ell}^\wedge \rightarrow (\mathrm{colim} K\mathrm{Rep}_{\mathcal{F}_i}\langle \mathcal{G}_i \rangle)_{\alpha_\ell}^\wedge.$$

Therefore it suffices to show that $(\mathrm{colim} K\mathcal{T}_i)_{\alpha_\ell}^\wedge$ is contractible.

Let $\mathcal{T} := \operatorname{colim} \mathcal{T}_i$, let $\mathcal{R}_i := \operatorname{Rep}_{k_i} \langle \mathcal{G}_i \rangle$, and let $\mathcal{R} := \operatorname{colim} \mathcal{R}_i$. The derived completion $(K\mathcal{T})_{\alpha_\ell}^\wedge$ is the cosimplicial $K\mathcal{R}$ -module whose n -simplices are obtained by $n+1$ applications of the functor $(\mathbb{H}_\ell \wedge_{K\mathcal{R}} -)$ to $K\mathcal{T}$. It suffices to show that $\mathbb{H}_\ell \wedge_{K\mathcal{R}} K\mathcal{T}$, which is isomorphic to $\operatorname{colim}(\mathbb{H}_\ell \wedge_{K\mathcal{R}_i} K\mathcal{T}_i)$, is contractible.

We will now show that for every i , the K -theory spectrum $K\mathcal{T}_i$ is weakly equivalent to $K\mathcal{R}_i$. Let $\mathbb{X} \subset \mathbb{N}$ be the subcategory of positive integers whose reciprocal occurs as an exponent of x in the ring $\mathcal{O}_i$. For every $s \in \mathbb{X}$ define $\mathcal{Z}_i(s) \subset \mathcal{T}_i$ to be the full subcategory of modules that are killed by $x^{1/s}$. Then by devissage [Qui73], the K -theory spectrum $K\mathcal{Z}_i(s)$ is weak equivalent to $K\mathcal{T}_i$. On the other hand, the category $\mathcal{Z}_i(s)$ is equivalent to the category $\operatorname{Rep}_{\mathcal{O}_i[x^{(1/s)}]} \langle \mathcal{G}_i \rangle$ via the forgetful functor.

Because for every positive integer r , the category $\mathcal{Z}_i(rs)$ is a full subcategory of $\mathcal{Z}_i(s)$, the functor $\mathcal{Z}_i(-)$ takes the category $\mathbb{X}$ contravariantly to the category of full subcategories of $\mathcal{T}_i$; it is therefore a cofiltered diagram. The category $\mathcal{R}_i$ maps via $(\mathcal{O}_i[x^{(1/s)}] \otimes_{k_i} -)$ to $\operatorname{Rep}_{\mathcal{O}_i[x^{(1/s)}]} \langle \mathcal{G}_i \rangle$, and it is easy to see that it is the limit

$$\mathcal{R}_i = \lim_{s \in \mathbb{X}} \operatorname{Rep}_{\mathcal{O}_i[x^{(1/sr)}]} \langle \mathcal{G}_i \rangle.$$

Applying the functor K yields the commutative diagram

$$\begin{array}{ccccccc} K\mathcal{Z}_i & \longrightarrow & \dots & \xrightarrow{\simeq} & K\mathcal{Z}_i(sr) & \xrightarrow{\simeq} & K\mathcal{Z}_i(s) \xrightarrow{\simeq} K\mathcal{T}_i \\ \downarrow \simeq & & & & \downarrow \simeq & & \downarrow \simeq \\ K\mathcal{R}_i & \longrightarrow & \dots & \longrightarrow & K\operatorname{Rep}_{\mathcal{O}_i[x^{(1/sr)}]} \langle \mathcal{G}_i \rangle & \longrightarrow & K\operatorname{Rep}_{\mathcal{O}_i[x^{(1/s)}]} \langle \mathcal{G}_i \rangle \end{array}$$

where $\mathcal{Z}_i$ is defined to be the limit $\lim_{s \in \mathbb{N}} \mathcal{Z}_i(s)$. This gives us the weak homotopy equivalence,

$$K\mathcal{T}_i \simeq K\mathcal{Z}_i \simeq K\mathcal{R}_i.$$

Now, since π_* commutes with filtered colimits and smash products, the above equivalence yields

$$(5) \quad \pi_*(\mathbb{H}_\ell \wedge_{K\mathcal{R}} K\mathcal{T}) \cong \operatorname{colim} \pi_*(\mathbb{H}_\ell \wedge_{K\mathcal{R}_i} K\mathcal{Z}_i) \cong \operatorname{colim} \pi_* \mathbb{H}_\ell.$$

Since $\pi_n \mathbb{H}_\ell = 0$ for all $n \neq 0$, we restrict our attention to π_0 . Because $\mathcal{Z}_i$ is closed under tensor product, $K\mathcal{Z}_i$ is actually a ring spectrum, so $\pi_0 K\mathcal{Z}_i$ is a ring, and

$$\pi_0(\mathbb{H}_\ell \wedge_{K\mathcal{R}_i} K\mathcal{Z}_i) = \pi_0 \mathbb{H}_\ell \otimes \pi_0 K\mathcal{Z}_i,$$

where the tensor product is taken over $\pi_0 K\mathcal{R}_i$. As a free $\pi_0 K\mathcal{R}_i$ -module, $\pi_0 K\mathcal{Z}_i$ is generated by a 1-dimensional k_i -vector space V_i that is annihilated by all (possibly fractional) powers of X in $\mathcal{O}_i$. Under τ_i^j , the module V_i (which corresponds to 1 in $\pi_0 \mathbb{H}_\ell$) gets sent to the module

$$V_i \otimes_{\mathcal{O}_i} \mathcal{O}_j = k_j [[x^{(1/\ell)}, x^{(1/p)}]] [x^{1/\ell^j}]$$

(which corresponds to ℓ^j in the ring $\pi_0 \mathbb{H}_\ell = \mathbb{F}_\ell$), and thus all of the maps in the colimit equation (5) are zero maps. Therefore $\pi_*(\mathbb{H}_\ell \wedge_{K\mathcal{R}} K\mathcal{T})$ is contractible.

The n -simplices of $(K\mathcal{T})_{\alpha_\ell}^\wedge$ are of the form $\mathbb{H}_\ell \wedge_{K\mathcal{R}} \dots \wedge_{K\mathcal{R}} \mathbb{H}_\ell \wedge_{K\mathcal{R}} K\mathcal{T}$ so they are all contractible as well. Since a level-wise weak equivalence of cosimplicial objects induces a weak equivalence on total spaces, $K\mathcal{T}_{\alpha_\ell}^\wedge$ is contractible, and thus

$$(\operatorname{colim} K\operatorname{Rep}_{\mathcal{O}_i} \langle \mathcal{G}_i \rangle)_{\alpha_\ell}^\wedge \xrightarrow{\simeq} (K\operatorname{Rep}_{\mathcal{T}^t} \langle \mathcal{G} \rangle)_{\alpha_\ell}^\wedge.$$

□

The following theorem follows immediately from the two preceding propositions.

Theorem 5.4. *Extension of scalars induces a weak equivalence of completed K -theory spectra,*

$$(K\mathrm{Rep}_k(\mathcal{G}))_{\alpha_\ell}^\wedge \xrightarrow{\cong} (K\mathrm{Rep}_{\mathcal{F}^t}(\mathcal{G}))_{\alpha_\ell}^\wedge \cong (K\mathcal{F})_{\alpha_\ell}^\wedge.$$

REFERENCES

- [AG60] M. Auslander and O. Goldman. The Brauer group of a commutative ring. *Trans. Amer. Math. Soc.*, 97:367–409, 1960.
- [BK72] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*, volume 304 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1972.
- [Car07a] G. Carlsson. Derived completions in homotopy theory. *Journal of Pure and Applied Algebra*, 2007. Submitted.
- [Car07b] G. Carlsson. Derived representation theory and the algebraic K -theory of fields. *Adv. Math.*, 2007. Submitted.
- [EKMM97] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [EM06] A. D. Elmendorf and M. A. Mandell. Rings, modules, and algebras in infinite loop space theory. *Adv. Math.*, 205(1):163–228, 2006.
- [LR07] G. Lyo and K. Ribet. Semisimple skew group rings and their modules, 2007.
- [May72] J. P. May. *The geometry of iterated loop spaces*. Springer-Verlag, Berlin, 1972. *Lectures Notes in Mathematics*, Vol. 271.
- [Qui73] D. Quillen. *Higher algebraic K-theory I*, volume 341 of *Lecture Notes in Mathematics*. Springer-Verlag, 1973.
- [Sus84] A.A. Suslin. On the K -theory of local fields. In *Proceedings of the Luminy conference on algebraic K-theory (Luminy, 1983)*, Vol. 2-3, pages 301–318. J. Pure Appl. Algebra, 1984.
- [Swa68] R. G. Swan. *Algebraic K-theory*, volume 76 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1968.

Department of Mathematics, Massachusetts Institute of Technology
 Cambridge, MA 02139
 gracelyo@math.mit.edu